Subject Code: R13103/R13

Set No - 1

I B.Tech I Semester Regular/Supple. Examinations Nov./Dec. - 2015 ENGINEERING PHYSICS

(Common to ECE, EEE, EIE, Bio-Tech, EComE, Agri.E)

Time: 3 hours Max. Marks: 70

Question Paper Consists of **Part-A** and **Part-B** Answering the question in **Part-A** is Compulsory, Three Questions should be answered from **Part-B**

PART-A

- 1. (a) Describe phenomenon of interference in thin films.
 - (b) Explain Bragg's law for X-ray diffraction.
 - (c) Mention various types of polarizations in dielectrics. and explain how they vary with temperature.
 - (d) State Gauss divergence theorem.
 - (e) What are matter waves and list out their properties?
 - (f) Distinguish between direct and indirect energy bandgap semiconductors.

[4+4+4+4+3+3]

PART-B

- 2. (a) Qualitatively analyse the spectrum obtained when a plane diffraction grating is exposed to monochromatic light of wavelength, λ .
 - (b) A grating has 6000 lines/cm. Find the angular separation between two wavelengths of 500nm and 510nm in the 3^{rd} order.
 - (c) Explain the concept of effective mass of a hole.

[8+4+4]

- 3. (a) Obtain an expression for acceptance angle of an optical fibre and express its numerical aperture in terms of fractional refractive index change.
 - (b) Calculate the numerical aperture of an optical fibre in air, if the refractive indices of its core and cladding are 2.5 and 2.495 respectively.
 - (c) Explain the principle of working of an LED.

[8+4+4]

- 4. (a) Derive the London equations and explain how their solution explains Meissner effect.
 - (b) The polarizability of ammonia molecule is found approximately by the measurement of dielectric constant as $2.42 \times 10^{-39} \, \text{C}^2 \, m \, / \, N$ and $1.74 \times 10^{-39} \, \text{C}^2 \, m \, / \, N$ at 309 K and 448 K respectively. Calculate the orientation polarizability at each temperature. (Given that $k_B = 1.38 \times 10^{-23} \, \text{J/K}$)
 - (c) Explain the phenomenon of double refraction.

[8+4+4]

- 5. (a) Discuss the factors affecting the architectural acoustics of a building and their remedies.
 - (b) A hall of volume 5500m³ is found to have a reverberation time of 2.3s. The sound absorbing surface of the hall has an area of 750m². Calculate the average absorption coefficient.
 - (c) Explain hysteresis loop of a ferromagnetic material.

[8+4+4]

WWW.MANARESULTS.CO.IN

- 6. (a) Derive an expression for density of energy states in metals.
 - (b) The density and atomic weight of copper are 8900 kg/m³ and 63.5. The relaxation time of electrons in Cu at 300K is 10⁻¹⁴ s. Calculate the electrical conductivity of copper.
 - (c) Express Maxwell's equations in differential form.

- 7. (a) What is Hall Effect? Deduce an expression for Hall coefficient.
 - (b) An n-type semiconducting specimen has a Hall coefficient of $3.66 \times 10^{-11} \text{m}^3/\text{As}$. The conductivity of the specimen is found to be $112 \times 10^7 \ \Omega^{-1} \text{m}^{-1}$. Calculate the charge carrier density and the electron mobility at room temperature.
 - (c) Distinguish between spontaneous and stimulated emissions.

[8+4+4]

Subject Code: R13103/R13

Set No - 2

I B. Tech I Semester Regular/Supple. Examinations Nov./Dec. - 2015 ENGINEERING PHYSICS

(Common to ECE, EEE, EIE, Bio-Tech, EComE, Agri.E)

Time: 3 hours Max. Marks: 70

Question Paper Consists of **Part-A** and **Part-B** Answering the question in **Part-A** is Compulsory, Three Questions should be answered from **Part-B**

PART-A

- 1. (a) Explain why the central fringe in Newton's rings is dark in the case of reflected system.
 - (b) How light waves are guided in an optical fibre? Explain.
 - (c) Distinguish between Type-I and Type-II superconductors
 - (d) State and explain Stoke's theorem.
 - (e) Explain the concept of the effective mass of an electron.
 - (f) Write short notes on drift and diffusion currents.

[4+4+4+4+3+3]

PART-B

- 2. (a) Explain Rayleigh's criterion for resolution. Obtain an expression for the resolving power of a diffraction grating.
 - (b) A grating of width 2inches is ruled with 15000 lines per inch. Find the smallest wavelength separation that can be resolved in second order at a mean wavelength of 500nm.
 - (c) Describe conductivity in an intrinsic semiconductor

[8+4+4]

- 3. (a) Explain the construction and working of He-Ne laser with energy level diagram. What are the merits of He-Ne laser?
 - (b) An optical fibre has a core of refractive index 1.5 and a cladding of refractive index 1.45. The diameter of the core of the fibre is 100µm and the medium surrounding the fibre is air. Determine (i) Numerical Aperture (ii) Acceptance angle
 - (c) Describe the principle behind working of a Photoconductor.

[8+4+4]

- 4. (a) Explain the phenomenon of superconductivity. Outline the BCS theory of superconductivity.
 - (b) Determine the critical current for a wire having diameter of 1mm at 4.2K. Critical temperature of the material is 8K and H_C at 0K is 6 x 10^4 A/m.
 - (c) Distinguish between a polarized light and an unpolarized light.

[8+4+4]

- 5. (a) Define sound absorption coefficient of a material and describe a method for its determination.
 - (b) A hall has a volume of 2265m³ and its total absorption is equivalent to 92.9m² of open window. What will be the effect on reverberation time if audience fill the hall and thereby increase the absorption by another 92.9m².
 - (c) Enumerate the properties of paramagnetic materials.

[8+4+4]

Page 1 of 2 WWW.MANARESULTS.CO.IN

- 6. (a) Explain Fermi-Dirac distribution function. Describe how the Fermi function varies with temperature.
 - (b) Find the relaxation time of conduction electrons in a metal of resistivity $1.54 \times 10^{-8} \Omega m$ if the metal has 5.8×10^{28} conduction electrons/m³.
 - (c) Express Maxwell's equations in integral form.

- 7. (a) Obtain an expression for carrier concentration in an intrinsic semiconductor.
 - (b) Calculate the intrinsic carrier concentration of Ge at 300K using the data E_g =0.7eV, and effective mass of electron and hole is $0.55m_e$ and $0.37m_e$ respectively.
 - (c) What are Miller indices? Explain.

[8+4+4]

I B. Tech I Semester Regular/Supple. Examinations Nov./Dec. - 2015 ENGINEERING PHYSICS

(Common to ECE, EEE, EIE, Bio-Tech, EComE, Agri.E)

Time: 3 hours Max. Marks: 70

Question Paper Consists of **Part-A** and **Part-B** Answering the question in **Part-A** is Compulsory, Three Questions should be answered from **Part-B**

PART-A

- 1. (a) Derive the expressions for thickness of quarter wave plate and half wave plate.
 - (b) Explain the terms (i) Basis (ii) Unit cell (iii) Crystal lattice and (iv) Lattice parameters
 - (c) Describe how ferromagnetic susceptibility varies with temperature.
 - (d) Define the term coefficient of absorption and write short notes on it.
 - (e) Outline the Sommerfeld's quantum free electron theory.
 - (f) Write short note on solar cells.

[4+4+4+3+3]

PART-B

- 2. (a) Derive an expression for the diameter of the nth dark ring in Newton's rings viewed under reflected system.
 - (b) In Newton's rings experiment the diameter of the 10th dark ring changes from 1.40cm to 1.27cm when a liquid is introduced between the lens and the glass plate. Calculate the refractive index of the liquid
 - (c) Explain the drift velocity and relaxation time of free electrons in metals.

[8+4+4]

- 3. (a) Describe the seven systems of crystals with suitable diagrams.
 - (b) Copper has FCC structure and its atomic radius is 0.1278nm. Calculate inter planar spacing for (111) and (321) planes.
 - (c) Explain Einstein's relation for mobility and diffusion coefficient of charge carriers.

[8+4+4]

- 4. (a) Define electronic polarization and polarizability. Deduce an expression for electronic polarizability in terms of radius of atom.
 - (b) Calculate the electronic polarizability of argon atom. Given that ε_r =1.0024 at NTP and N=2.7x10²⁵ atoms/m³.
 - (c) How do you determine the refractive index of a liquid using Newton's rings experiment? [8+4+4]
- 5. (a) State and explain Sabine's formula for reverberation time of a hall. Derive Sabine's formula for reverberation time.
 - (b) A hall has dimensions 20 x 15 x 5m³. The reverberation time is 3.5s. Calculate the total absorption of its surface and the average absorption coefficient.
 - (c) Describe the FCC crystal structure.

[8+4+4]

- 6. (a) How does the band theory of solids lead to the classification of solids into conductors, semiconductors and insulators?
 - (b) Calculate the drift velocity of conduction electrons in copper at a temperature of 300K when a copper wire of length 2m and resistance 0.02Ω carries a current of 15A. Given that mobility of the electrons is $4.3 \times 10^{-3} \, \text{m}^2/\text{Vs}$.
 - (c) How does a SQUID work?

- 7. (a) Derive an expression for carrier concentration in an n-type semiconductor.
 - (b) The energy gap of Si is 1.1eV. The average electron effective mass is $0.31m_e$, where m_e is the free electron mass. Calculate the concentration of electrons in the conduction band of Si at room temperature, T=300K. Assume that $E_F = E_g/2$.
 - (c) Explain in detail the acoustic demands of a hall.

[8+4+4]

Subject Code: R13103/R13

Set No - 4

I B. Tech I Semester Regular/Supple. Examinations Nov./Dec. - 2015 ENGINEERING PHYSICS

(Common to ECE, EEE, EIE, Bio-Tech, EComE, Agri.E)

Time: 3 hours Max. Marks: 70

Question Paper Consists of **Part-A** and **Part-B** Answering the question in **Part-A** is Compulsory, Three Questions should be answered from **Part-B**

PART-A

- 1. (a) Discuss in detail the phenomenon of double refraction.
 - (b) Explain how X-rays can be made to diffract?
 - (c) Explain Meissner effect in superconductors.
 - (d) Explain the terms 'Reverberation' and 'Reverberation time'.
 - (e) Define relaxation time and mobility of charge carriers.
 - (f) Describe the working of an LED.

[4+4+4+4+3+3]

PART-B

- 2. (a) Analyze qualitatively Fraunhofer diffraction at double slit with suitable diagrams.
 - (b) A plane transmission grating with 5000 lines/cm gives a second order diffraction maximum at an angle of 30⁰ from the central maximum. Find (i) the wavelength of light diffracted and (ii) the maximum order of diffraction possible.
 - (c) Write down any four applications of Hall effect.

[8+4+4]

- 3. (a) Define the terms coordination number, atomic radius and packing density. Calculate these factors for simple cubic, body centered cubic and face centered cubic crystals.
 - (b) A beam of X-rays of wavelength 0.071nm is diffracted by (110) plane of rock salt with lattice constant of 0.28nm. Find the glancing angle for the second order diffraction.
 - (c) What are the drawbacks of classical free electron theory?

[8+4+4]

- 4. (a) Distinguish between ferro, anti-ferro and ferri magnetic materials in terms of susceptibility and its dependence on temperature.
 - (b) An electron is moving in a circular orbit of radius 0.62 x 10⁻¹⁰m. The electron performs 10¹⁵ revolutions per second. Determine the magnetic moment associated with the orbital motion of the electron.
 - (c) State and explain Eyring's formula.

[8+4+4]

- 5. (a) By using Gauss divergence and Stokes theorems convert Maxwell's equations from differential form to integral form.
 - (b) Explain lasing action in a three level system.
 - (c) Deduce the Claussius-Mossotti relation for dielectrics.

[8+4+4]

Page 1 of 2

WWW.MANARESULTS.CO.IN

- 6. (a) Discuss the Kronig-Penny model for the motion of an electron in a periodic potential.
 - (b) Use the Fermi Dirac distribution function to obtain the value of F(E) for E-E_F=0.01eV at 200K. Given that $k_B = 1.38 \times 10^{-23} J/K$.
 - (c) Outline the principle behind the working of an optical fibre.

- 7. (a) Derive an expression for carrier concentration in a p-type semiconductor.
 - (b) The forbidden gap in pure silicon is 1.1eV. Compare the number of conduction electrons at temperatures 37°C and 27°C.
 - (c) What is a diffraction grating? Explain with the help of a diagram.

[8+4+4]