

B.Tech II Year II Semester (R13) Supplementary Examinations May/June 2017 MATHEMATICS – II

(Computer Science and Engineering)

Max. Marks: 70

Time: 3 hours

1

PART – A

(Compulsory Question)

- Answer the following: (10 X 02 = 20 Marks)
 - (a) Determine the rank of the matrix: $\begin{bmatrix} 1 & 2 & 3 \\ 1 & 4 & 2 \end{bmatrix}$
 - (b) Find the values of k for which the system of equations: (3k-8)x + 3y + 3z = 0 3x + (3k-8)y + 3z = 0
 - 3x + (3x 8)y + 3z = 03x + 3y + (3k - 8)z = 0 has a nontrivial solution.
 - (c) Given the values.

X:	5	7	11	13	17
f(x)	150	392	1492	2366	5202

Evaluate f(9), using Lagrange's interpolation formula.

- (d) Use Simpson's $1/3^{rd}$ rule to find $\int_0^{0.6} e^{-x^2} dx$ by taking seven ordinates.
- (e) Apply Runge-Kutta fourth order method, to find an approximate value of y when x = 0.2, given that dy/dx = x+y and y = 1 when x = 0.
- (f) Find the Fourier series to represent x^2 in the interval (-I, I)
- (g) Find the Fourier cosine transform of e^{-x^2} .
- (h) Find the z-transform of the following:

(i) $3n = 4 \sin n\pi/4 + 5a$. (ii) $(n + 1)^2$.

- (i) Derive a partial differential equation (by eliminating the constants) from the equation $2z = \frac{x^2}{x^2} + \frac{y^2}{x^2}$.
- (j) Using the method of separation of variables solve $py^3 + qx^2 = 0$.

PART – B

(Answer all five units, $5 \times 10 = 50$ Marks)

- 2 (a) Find the matrix p which transforms the matrix $A = \begin{pmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{pmatrix}$ to the diagonal form. Hence calculate A⁴.
 - (b) Reduce the quadratic form 2xy + 2yz + 2zx into canonical form.

OF

- 3 (a) Prove that the matrix $A = \begin{pmatrix} \frac{1}{2}(1+i) & \frac{1}{2}(-1+i) \\ \frac{1}{2}(1+i) & \frac{1}{2}(1-i) \end{pmatrix}$ is unitary and find A⁻¹.
 - (b) Prove that every Hermitian matrix can be written as A + iB, where A is real and symmetric and B is real and skew-symmetric.

Contd. in page 2

www.ManaResults.co.in



UNIT - II

- (a) Find a real root of the equation $x^3 2x 5 = 0$ by the method of false position corrected to three 4 decimal places.
 - (b) Find the positive root of $x^4 x = 10$ corrected to three decimal places, using Newton-Raphson method.

(a) From the following table, estimate the number of students who obtained marks between 40 and 45: 5

Marks:	30-40	40-50	50-60	60-70	70-80	
No. of students:	31	42	51	35	31	

- (b) Evaluate $\int_0^6 \frac{dx}{1+x^2}$ by using:
 - (i) Trapezoidal rule.
 - (ii) Simpson's 1/3 rule.
 - (iii) Simpson's 3/8 rule.

UNIT - III)

- 6
- (a) Solve by Taylor's series method the equation $\frac{dy}{dx} = \log (xy)$ for y(1.1) and y(1.2), given y(1) = 2. (b) Apply Milne's method, to find a solution of the differential equation y' =x y² in the range $0 \le x \le 1$ for the boundary conditions y = 0 of x = 0.

OR

(a) Find the Fourier series to represent x - x^2 from x = - π to x = π . 7

(b) Express f(x) = x as a half range sine series in 0 < x < 2.

Find the Fourier transform of: $f(x) = \begin{cases} 1 - x^2, |x| \le 1\\ 0, & |x| > 1 \end{cases}$ 8 (a)

Hence evaluate
$$\int_0^\infty \frac{x \cos x - \sin x}{x^3} \cos \frac{x}{2} dx$$

(b) Show that $Z(\sin hn\theta) = \frac{1}{z^2 - 2z \cos h\theta + 1}$

OR

- (a) If $Z^{-1}[U(z)] = u_n$ and $Z^{-1}[V(z)] = v_n$, then prove that $Z^{-1}[U(z), V(z)] = \sum_{m=0}^n u_m$. $v_{n-m} = u_n * v_n$ where the symbol * denotes the convolution operation. 9
 - (b) Find the inverse z.-transform of $\frac{z^3-20z}{(z-2)^3(z-4)}$.

UNIT - V

- Solve the differential equation $\frac{\partial u}{\partial t} = \alpha^2 \frac{\partial^2 u}{\partial x^2}$ for the conduction of neat along a rod without radiation, 10 subject to the following conditions:
 - U is not infinite for $t \rightarrow \infty$. (a)

(b)
$$\frac{\partial u}{\partial x} = 0$$
 for $x = 0$ and $x = l$.

 $u = lx - x^2$ for t = 0, between x = 0 and x = l. (c)

OR

Solve the Laplace equation $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$ subject to the conditions u(0,y) = u(l,y) = u(x,0) = 011 and $u(x, a) = \sin n\pi x/l$.

www.ManaResults.co.in