

**Code No: 133BD****JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY HYDERABAD****B.Tech II Year I Semester Examinations, May/June - 2019****MATHEMATICS – IV****(Common to CE, EEE, ME, ECE, CSE, EIE, IT MCT, ETM MMT, AE, MIE, PTM, CEE, MSNT)****Time: 3 Hours****Max. Marks: 75****Note:** This question paper contains two parts A and B.

Part A is compulsory which carries 25 marks. Answer all questions in Part A.

Part B consists of 5 Units. Answer any one full question from each unit.

Each question carries 10 marks and may have a, b, c as sub questions.

**PART- A****(25 Marks)**

- 1.a) State the necessary and sufficient conditions for a function  $f(z) = u + iv$  to be analytic. [2]
- b) Show that  $f(z) = |z|^2$  is not analytic at any point. [3]
- c) State Cauchy's integral theorem. [2]
- d) Find the poles and the residues at the poles of the function  $f(z) = \frac{e^z}{\cos \pi z}$ . [3]
- e) Define bilinear transformation and cross ratio. [2]
- f) Find the image of the circle  $|z| = 2$ , under the transformation  $w = z + 3 + 2i$ . [3]
- g) State Fourier integral theorem. [2]
- h) Expand  $f(x) = \pi x - x^2$  in a half range sine series in  $(0, \pi)$ . [3]
- i) Classify the partial differential equation  $u_{xx} + 6u_{xy} + 2u_{yy} + 2u_x - 2u_y + u = x^2y$ . [2]
- j) Write the three possible solutions of the heat equation.  

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2}$$
 [3]

**PART-B****(50 Marks)**

- 2.a) If  $f(z)$  is a regular function of  $z$ , prove that  

$$\left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) |f(z)|^2 = 4 |f'(z)|^2.$$
  - b) Let  $f(z) = u(r, \theta) + iv(r, \theta)$  be an analytic function. If  $u = -r^3 \sin 3\theta$ , then construct the corresponding analytic function  $f(z)$  in terms of  $z$ . [5+5]
- OR**
- 3.a) Show that the function  $f(z)$  defined by  

$$f(z) = \frac{x^2 y^3 (x+iy)}{x^6 + y^{10}} \quad \text{for } z \neq 0$$
is not analytic at the origin, even though it satisfies the  
 $f(0) = 0$   
 Cauchy-Riemann equations at the origin.
  - b) Determine the analytic function whose real part is  $\log \sqrt{x^2 + y^2}$ . [5+5]

4. Represent the function  $\frac{1}{z^2-4z+3}$  in the domain  
 (a)  $1 < |z| < 3$  (b)  $|z| < 1$ . [10]

**OR**

- 5.a) Expand the function  $f(z) = \frac{z}{(z+1)(z+2)}$  about  $z = -2$ , and name the series thus obtained.

- b) Evaluate  $\oint_C \frac{e^z}{(z+3)(z+2)} dz$ , where  $C$  is the circle  $|z - 1| = \frac{1}{2}$ . [5+5]

6. Evaluate the integral using contour integration  $\int_0^{2\pi} \frac{d\theta}{2+\cos \theta}$ . [10]

**OR**

7. Show that the transformation  $w = i \frac{1-z}{1+z}$  transforms the circle  $|z| = 1$  into the real axis of  $w$  plane and the interior of the circle  $|z| < 1$  into the upper half of the  $w$  plane. [10]

8. Find the Fourier transform of  $f(x) = \begin{cases} 1-x^2, & \text{if } |x| < 1 \\ 0, & \text{if } |x| > 1. \end{cases}$  Hence evaluate

$$\int_0^\infty \frac{x \cos \frac{x-\sin x}{x^3} \cos \frac{x}{2}}{x^3} dx. \quad [10]$$

**OR**

- 9.a) Obtain the half range cosine series for

$$f(x) = \begin{cases} kx, & \text{for } 0 \leq x < L/2 \\ k(L-x), & \text{for } L/2 \leq x \leq L. \end{cases}$$

- b) Find the Fourier sine transform of  $f(x) = e^{-|x|}$ . Hence show that  $\int_0^\infty \frac{x \sin mx}{x^2+1} dx = \frac{\pi}{2} e^{-m}$ . [5+5]

10. A string is stretched and fastened to two points  $L$  apart. Motion is started by displacing the string in the form  $y = a \sin \frac{\pi x}{L}$  from which it is released at time  $t = 0$ . Find the displacement of any point at a distance  $x$  from one end at time  $t$ . [10]

**OR**

11. Write down the one dimensional heat equation. Find the temperature  $u(x, t)$  in a slab whose ends  $x = 0$  and  $x = L$  are kept at zero temperature and whose initial temperature  $f(x)$  is given by

$$f(x) = \begin{cases} k, & \text{when } 0 < x < \frac{1}{2}L \\ 0, & \text{when } \frac{1}{2}L < x < L. \end{cases} \quad [10]$$

---oo0oo---